arXiv:1612.09036v1 [math.RA] 24 Dec 2016

$\mathbb{F}_p$ and Z_p Valued Holomorphic Functions over Graphs

H. Mohades
Amirkabir University of Technology, Iran

December 30, 2016

Abstract

The definition of a holomorphic function over a general measurable space S endowed with a Markov process is defined by Zeghib and Barre. In this article we consider holomorphic functions over graphs whose ranges are a given finite field or a cyclic group. Also we consider a relation between $\mathbb{C}$-holomorphic functions over regular trees and the field of p-adic numbers.

AMS subject Classification 2010: 05-XX, 11Txx

Keywords: Finite field , Harmonic function, Holomorphic function, Discrete Laplacian Operator,

1 Introduction

Let R be a commutative ring. The definition of an R-holomorphic function over measurable space S endowed by a Markov process is defined in [1]. In the sense of [1] a function $\phi : S \to R$ is holomorphic if both ϕ and ϕ^2 are harmonic with respect to the natural notion of Laplacian over S. In the case of Riemannian manifolds this notion of $\mathbb{C}$-holomorphic functions coincides with the concept of harmonic morphism from a manifold to $\mathbb{R}^2$. Harmonic morphism from a geometric graph to $\mathbb{C}$ are constant maps. Nevertheless there are nonconstant $\mathbb{C}$-holomorphic function over graphs and therefore the definition of $\mathbb{C}$-holomorphic functions is different from the notion of holomorphic function from a graph to $\mathbb{C}$. A harmonic function on a graph or a Riemannian manifold can be constructed by going to infinity (martin boundary) and then reproducing the object using a Poisson integral. Versus this global way of producing a holomorphic function, the construction of an R-holomorphic function will be done locally. In fact, there is a "special dynamics" which allows one to construct the holomorphic functions. We show this phenomenon with an example on 3-regular graphs. The main part of this article is devoted to $\mathbb{F}_p$ or Z_p-valued holomorphic functions. In the case of $\mathbb{F}_p$ we provide a formula for the asymptotic behavior of the cardinality of the restriction of holomorphic functions to a neighborhood of a point in some regular trees. At the end of the article we determine a relation between $\mathbb{C}$-valued holomorphic functions and the field of p-adic numbers. In fact we prove that a subgroup of automorphisms of this field is the same as special set of $\mathbb{C}$-valued holomorphic functions on a $(p+1)$-regular tree.

2 Harmonic and holomorphic functions

A graph is an ordered pair $X = (V, E)$ of disjoint sets V and E with a map $o : E \to V^2$, where $V^2 = \{\{a, b\} | a \in V \text{ and } b \in V\}$. Two elements $e \in V$ and $e' \in V$ are adjacent if $o^{-1}\{e, e'\}$ is not

an empty set. For an element $e \in V$ let $N(e) = \{e' \in V | e' \sim e\}$, where $e \sim e'$ means e is adjacent to e'. Let $C(X, F)$ be the set of all F-valued functions on the set V. The elements of $C(X, F)$ will be called the functions over the graph X.

Definition 1. Let $f \in C(X, F)$, the Laplacian Δf is the function:

$$\Delta f(x) = \sum_{x \sim y} f(y) - f(x) \tag{1}$$

Definition 2. A function $g : X \to \mathbb{F}$ is called harmonic if $\Delta(g) = 0$. The set of harmonic functions is a vector subspace of the set of all $\mathbb{F}$-valued functions on X.

The only $\mathbb{R}$-valued($\mathbb{C}$-valued) harmonic functions on a finite graph are constant functions. Despite of this fact on $\mathbb{R}$-holomorphic functions, generally the set of harmonic F-valued functions are not constant on a finite graph.

Example 1. *Let X be a cyclic graph with 6 vertices and let $F = Z_3$. The set of Z_3-valued harmonic functions on X has nonconstant functions.*

Definition 3. Let $\phi : X \to \mathbb{F}$, where X is a Riemannian manifold or a graph. We say that ϕ is holomorphic if both ϕ and its square ϕ^2 are harmonic.

Example 2. *Let X be a cyclic graph, then the set of $\mathbb{F}_q$-holomorphic functions on X (q is an odd number) is the same as the set of constant functions.*

3 Holomorphic functions on finite fields

For the sake of simplicity we neglect finite fields of characteristic 2.

Theorem 1. *Let ϕ be a $\mathbb{F}_q$-valued holomorphic function over X then we have the following equations at each vertex.*

$$\sum_i a_i^2 = \sum_i a_i = 0 \tag{2}$$

when $a_i = \begin{cases} \phi(e_i) - \phi(e_0) & if \quad e_i \sim e_0 \\ 0 & otherwise \end{cases}$

Proof. By the definition of a holomorphic function we have

$$\Delta(\phi)(e_0) = \sum_{e_i \sim e_0} \phi(e_i) - \phi(e_0) = \sum_i a_i = 0$$

and

$$\Delta(\phi^2)(e_0) = \sum_{e_i \sim e_0} (\phi^2(e_i) - \phi^2(e_0)) = \sum_i (\phi(e_i)^2 + \phi(e_0)^2) - 2\phi(e_0) \sum_i \phi(e_i) =$$

$$\sum_i (\phi(e_i) - \phi(e_0))^2 = \sum_i a_i^2 = 0.$$

□

Theorem 2. *There are nontrivial $\mathbb{F}_q$-valued holomorphic functions on a tree with no vertex of degree less then 4.*

Proof. By the extension of the Chevalley's theorem we know that a set of equations consisting of a quadratic form and a linear one on more than 4 variables has nontrivial solutions on $\mathbb{F}_p$. So if we let $\phi(e_0) = c$, then we can find values for ϕ on $N(e_0)$ which are not equal with c. Now we extend this to the points which are joint to e_0 by a path of length 2. If $\phi(e_1) = d$ then we have the following set of equations at the vertex e_1.

$$\begin{cases} (c-d)^2 + \sum_{i=1}^{deg(e_1)-1}(y_i - d)^2 = 0 \\ (c-d) + \sum_{i=1}^{deg(e_1)-1}(y_i - d) = 0 \end{cases}$$

Let $y_i - d = x_i$. Now, we obtain $\sum_{i=1}^{deg(e_1)-2} x_i^2 = b$ for some $b \in \mathbb{F}_p$. We know that the equation

$$x_1^2 + x_2^2 = b$$

has at least $p-1$ solutions [4]. Considering $deg(e_1)-2 \geq 2$, we are able to extend ϕ to a holomorphic function. □

Theorem 3. *Let $Hol_{e_0,s}(\mathbb{F}_q)$ denotes the restriction of the set of $\mathbb{F}_p$-valued holomorphic functions ϕ such that $\phi(e_0) = s$, to $N(e_0)$. Then $card(Hol_{e_0,s}(\mathbb{F}_q)) = q^{deg(e_0)-2} + (q-1)q^{\frac{deg(e_0)-3}{2}}\eta$ when $\eta \in \{1, -1\}$.*

Proof. From the equation 2 it is enough to find the number of solutions of the equation

$$\sum_{1<i<j,j=2}^{deg(e_0)} (a_j - a_i)^2 = 0$$

The number of solution of this equation is equal with the number of solutions of the nondegenerated diagonal quadratic equation in $deg(e_i) - 1$ variables. Now the theorem is a result of Lebesgue theorem on the counting of number of solutions of a quadratic equation. □

Now we provide some facts on the cardinality of the restriction of holomorphic functions to a finite neighborhood of a vertex.

Theorem 4. *Let G be a tree and p divides the degree of each vertex when p is the characteristic of the field $\mathbb{F}_q$.Let $Hol_{e_0,e_1,s,u}(\mathbb{F}_q)$ denotes the set of $\mathbb{F}_q$ valued holomorphic functions ϕ such that $\phi(e_0) = s, \phi(e_1) = u$, restricted to the set $N(e_1)$. Then $card(Hol_{e_0,e_1,s,u}(\mathbb{F}_p)) = q^{p-3}$.*

Proof. For $e_j \sim e_1 \quad , j = 1, ..., p-1$, let $x_j = \phi(e_j) - u$, also let $t = u - s$. Then, we have the following set of equations.

$$\begin{cases} \sum_{j=1}^{p-1} x_j^2 = -t^2 \\ \sum_{j=1}^{p-1} x_j = -t \end{cases}$$

Now we are in the set up of [4] (page 341). Where $b = \sum_j b_j^2 a_j^{-1} = p-1$, and $c = b_0^2 - a_0 b = (-t)^2 - (p-1)(-t^2) = pt^2 = 0$. The number of variables is equal with $p-1$ which is an even number so the cardinality of solutions is q^{p-3}. □

Corollary 1. *In a p-regular graph the cardinality of the set of functions obtained by the restriction of $\mathbb{F}_q$-valued holomorphic functions to an r-neighborhood of e_0 is equal with*

$$[q^{p-2} + (q-1)q^{\frac{p-3}{2}}\eta]q^{(p-3)(r-1)}.$$

Proof. The proof is an straightforward application of theorems 3 and 4. □

Corollary 2. *In a 3-regular tree(Tr_3), there are only finitely many $\mathbb{F}_{3^n}$-valued holomorphic functions up to permutation.*

Proof. It is a result of Theorem 4. □

Theorem 5. *Let L be a finite graph with n vertices and $|E|$ edges. Let $|E| > 3n$ ($\chi(L) > 2n$). Then there are nontrivial $\mathbb{F}_q$-valued holomorphic functions over L.*

Proof. By the Theorem 2 we must solve a system of $2n$ equations on E variables. n equations have a degree equal with 2 and the other equations have degree equal with 1. Summation of equation's degree are equal with $3n$. Therefore, by Theorem 6.8. of [4] there exists nontrivial solutions for this set of equations. So, we have nontrivial holomorphic functions on these finite graphs. □

4 Z_p-valued Holomorphic functions and dynamics

Theorem 6. *There are nontrivial Z_p $p > 5$ valued holomorphic functions on a tree with no vertex of degree less than 6.*

Proof. The first part of the proof is similar to the proof of theorem 2. For the last part of the discussion we must prove that the equation $\sum_{i=1}^{n} x_i^2 = b(mod\ p)$ for $n > 4$ has a nontrivial solution. Now we use the Lagrange's four-square theorem [3] which asserts that every natural number is equal with the summation of at most 4 square numbers. Therefore every choice of fifth variable leads to a solution which is not trivial. □

Regular trees are important in the study of p-adic field $\mathbb{Q}_p$ and p-adic ring $\mathbb{Z}_p$ which is the inverse limit of the finite rings Z_{p^n}. The Bruhat-Tits tree of a p-adic field is a regular $p+1$- tree.

Let s, t be two classes of neighboring elements of the Bruhat-Tits tree over $\mathbb{Q}_2$. This tree is isomorphic to Tr_3. Then we have the following Theorem.

Theorem 7. *Let $Aut_{s,t}(\mathbb{Q}_2)$ be the set of automorphisms of the $2-$adic field $\mathbb{Q}_2$ which preserve s and t. Then there is a one-one correspondence between $Aut_{s,t}(\mathbb{Q}_2)$ and $\mathbb{C}$-valued holomorphic functions over the Bruhat-tits tree which have fixed values α and β ,$\alpha \neq \beta$, on s and t.*

Proof. Let A_1 and A_2 be two connected subtrees of Tr_3 obtained by deleting the edge $\bar{st}$. Let $A(s,t) = A_1 \cup \bar{st}$, $A(t,s) = A_2 \cup \bar{st}$. Let $H_{\alpha,\beta}$ be the space of holomorphic functions on $A(s,t)$ with prescribed values $\phi(s) = \alpha$ and $\phi(t) = \beta$. Let $Aut(A)$ be the automorphism group of the graph $A(s,t)$. s and t do not change under the action of the elements of $Aut(A)$. Also, for any element w of $A(s,t)$, there are elements of the group which exchange two neighborhoods of w which are not between w and s. On the other hand, this exchanging was the only freedom in determining the holomorphic function(which is obtained by the induced dynamics to choose j or j^2 [1]). Equivalently, right action of $Aut(A(s,t))$ on $H_{\alpha,\beta}$ is transitive. Now it is easy to see that the action of $Aut(A(s,t)) \times Aut(A(t,s))$ on $\mathbb{C}$-valued holomorphic functions over the Bruhat-tits tree with the fixed values $\alpha \neq \beta$ on s, t is simply transitive. This completes the proof. □

Let we consider the special case of $G = Tr_3$. There is no Z_3-holomorphic function on $G = Tr_3$. Now we consider the set of Holomorphic functions from G to Z_9. The equation $a + b + c = a^2 + b^2 + c^2 = 0$ has 3 different solutions(up to permutation) $(0,0,0), (3,3,3), (0,3,6)$ so we can

construct the set of holomorphic functions using a dynamics over G. In fact let we consider the dynamics obtained by the random choice of elements 0, 3 or 6 on edge initiated from each vertex. This dynamics is equivalent to all holomorphic functions on Tr_3 when we neglect the neighboring edges with constant values 0 or 3.

References

[1] S. Barre, A. Zeghib, Real and discrete holomorphy: Introduction to an algebraic approach, J. Math. Pures Appl. 87 (2007).

[2] M. Hubenthal, Maximal subspsces of zeros of quadratic forms over finite fields, https://www.whitman.edu/Documents/Academics/Mathematics/hubentjm.pdf, (2006).

[3] K. Ireland, M. Rosen, A classical introduction to modern number theory, Springer, (1990).

[4] R. Lidl, H. Niederreiter, P.M. Cohn, Finite fields, cambridge university press , 1997.

[5] p-adic numbers, www1.spms.ntu.edu.sg/ frederique/antchap5.pdf.

[6] A. Weil, Numbers of solutions of equations in finite fields, Bull. Amer. Math. Soc. 55 (1949).

[7] j. Igusa, An Introduction to the Theory of Local Zeta Functions, AMS International Press,(2007)

[8] A.V. Zabrodin, Non-Archimedean Strings and Bruhat-Tits Trees, Springer-Verlag, (1989).

[9] S. R. Ghorpade, G. Lachaud, Number of solutions of equations over finite fields and a conjecture of Lang and Weil, Number Theory and Discrete Mathematics (Chandigarh, 2000).

[10] D.Wan, Computing Zeta Functions Over Finite Fields, Contemporary Mathematics,(1998).